


KI Administration Based on FDA (2001)
 (Thyroid CDE from Radioiodine)



Map Size: 9.2 km by 9.2 km Id: Beta.rcE17326.rcC1
 Approved by: {NARAC Operations; NARAC}

Early Phase Dose and Actions

Description	(rem) Extent Area	Population
Adult thyroid CDE - Early Phase FDA Guidance for KI administration to adults over 40.	>500 0.2km 0.008 km2	10
Adult thyroid CDE - Early Phase FDA Guidance for KI administration to adults under 40.	>10 5.8km 12.7 km2	30,500
Adult thyroid CDE - Early Phase FDA Guidance for KI administration to pregnant or lactating females.	>5 6.8km 19.6 km2	39,500
Child thyroid CDE - Early Phase PAG for KI administration to children under 18.	>5 8.0km 33.1 km2	64,000

Areas and counts in the table are cumulative. Population Source = LandScanUSA10NARAC9.

Effects or contamination from June 30, 2011 13:00 UTC to July 04, 2011 13:00 UTC
 Release Location: 36.715775 N, 121.623420 W
 Material: I-133
 Generated On: October 17, 2012 18:43 UTC
 Model: ADAPT/LODI
 Comments: Doses shown are total accumulated from the beginning of release.
 Hypothetical release